

BEST AVAILABLE COPY



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁵ : A61K 39/00, 39/02, C12N 1/10, 1/36</p>	A1	<p>(11) International Publication Number: WO 94/16725</p> <p>(43) International Publication Date: 4 August 1994 (04.08.94)</p>		
<table style="width: 100%; border: none;"> <tr> <td style="width: 50%; vertical-align: top; padding: 5px;"> <p>(21) International Application Number: PCT/US94/00075</p> <p>(22) International Filing Date: 3 January 1994 (03.01.94)</p> <p>(30) Priority Data: 006,349 19 January 1993 (19.01.93) US</p> <p>(60) Parent Application or Grant (63) Related by Continuation US 006,349 (CON) Filed on 19 January 1993 (19.01.93)</p> <p>(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): SCHMATZ, Dennis, M. [US/US]; 306 Elizabeth Avenue, Cranford, NJ 07016 (US). COX, James, L. [US/US]; 536 Lyme Rock Road, Bridgewater, NJ 08807 (US).</p> <p>(74) Agent: ROSE, David, L.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p> </td> <td style="width: 50%; vertical-align: top; padding: 5px;"> <p>(81) Designated States: AU, BB, BG, BR, BY, CA, CN, CZ, FI, HU, JP, KR, KZ, LK, LV, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, UZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p> </td> </tr> </table>			<p>(21) International Application Number: PCT/US94/00075</p> <p>(22) International Filing Date: 3 January 1994 (03.01.94)</p> <p>(30) Priority Data: 006,349 19 January 1993 (19.01.93) US</p> <p>(60) Parent Application or Grant (63) Related by Continuation US 006,349 (CON) Filed on 19 January 1993 (19.01.93)</p> <p>(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): SCHMATZ, Dennis, M. [US/US]; 306 Elizabeth Avenue, Cranford, NJ 07016 (US). COX, James, L. [US/US]; 536 Lyme Rock Road, Bridgewater, NJ 08807 (US).</p> <p>(74) Agent: ROSE, David, L.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p>	<p>(81) Designated States: AU, BB, BG, BR, BY, CA, CN, CZ, FI, HU, JP, KR, KZ, LK, LV, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, UZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p>
<p>(21) International Application Number: PCT/US94/00075</p> <p>(22) International Filing Date: 3 January 1994 (03.01.94)</p> <p>(30) Priority Data: 006,349 19 January 1993 (19.01.93) US</p> <p>(60) Parent Application or Grant (63) Related by Continuation US 006,349 (CON) Filed on 19 January 1993 (19.01.93)</p> <p>(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): SCHMATZ, Dennis, M. [US/US]; 306 Elizabeth Avenue, Cranford, NJ 07016 (US). COX, James, L. [US/US]; 536 Lyme Rock Road, Bridgewater, NJ 08807 (US).</p> <p>(74) Agent: ROSE, David, L.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p>	<p>(81) Designated States: AU, BB, BG, BR, BY, CA, CN, CZ, FI, HU, JP, KR, KZ, LK, LV, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, US, UZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p>			
<p>(54) Title: LIVE COCCIDIOSIS VACCINE</p> <p>(57) Abstract</p> <p>Live sporulated oocysts are administered to one day old chickens to provide immunity against coccidiosis without the need to provide supplemental anticoccidial therapy.</p>				

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

TITLE OF THE INVENTION
LIVE COCCIDIOSIS VACCINE

BACKGROUND OF THE INVENTION

5 It is known that infective stages of coccidial organisms, such as sporulated oocysts, can be administered to neonatal fowl to induce immunity against coccidiosis. However, administration of such "wild-type" organisms must be followed by the administration of anticoccidial agents to prevent the outbreak of coccidiosis by the infective organisms. In addition, the administration of such infective strains requires that the fowl, preferably chickens, be maintained on litter to provide for a constant level of reinfection to stimulate lasting immunity. This technique also suffers failure when too many or too few oocysts are administered. Being infective, too many oocysts can produce an actual outbreak of coccidiosis while too few oocysts result in inadequate levels of immunization leaving the chickens susceptible to natural outbreaks of coccidiosis.

 One attempt at avoiding the problems associated with live vaccines is to administer attenuated precocious strains of Eimeria. These strains are less infective than the "wild-type" and tend to avoid problems associated with vaccines based on such strains. "Attenuated" strains are those that have been serially passaged through the host or through eggs, or a combination of host and egg passages, so that the resultant strain is less infective than the parent strain. "precocious" strains are those that have been developed to have short prepatent periods, which is the interval between the ingestion of an infective stage and its development within the host leading to the excretion of the next infective stage of the parasite. Attenuated strains are capable of conferring immunity and precocious strains are capable of developing into the next stage, however, the reduced level of infectivity of the attenuated strains, combined with the shortened development period of the parasite induces immunity without pathology and thus the host will not suffer ill effects from the vaccination.

- 2 -

However, such attenuated precocious strains, because they are less infective must be administered at higher levels, based on the number of oocysts administered, and it is recommended that vaccination with sporulated oocysts not take place earlier than three days after birth. See U.S. Patents 4,438,097 to Shirley and 5,055,292 to McDonald, et al.

SUMMARY OF THE INVENTION

The instant invention provides for the full immunization of fowl against coccidiosis by the oral administration of live sporulated oocysts of attenuated precocious strains of Eimeria at day one of age without the subsequent administration of anticoccidial agents. Thus, it is an object of this invention to describe the administration of such live, infective, but attenuated and precocious oocysts to newly hatched fowl. It is a further object to describe the effects such administration has upon the ability of such fowl to withstand a challenge of coccidiosis. A still further object is to describe formulations and methods to administer the oocysts to such fowl. Further objects will become apparent from a reading of the following description.

DESCRIPTION OF THE INVENTION

This invention is concerned with the administration of live, attenuated, precocious strains of coccidial organisms, in particular strains of Eimeria, to newly hatched fowl, in particular, chickens, but also turkey, ducks, quail and the like. The vaccine may be univalent, with a single species of Eimeria in the vaccine, but preferably a polyvalent vaccine is employed containing from two to all seven of the common species of Eimeria, viz E. necatrix, E. aservulina, E. brunetti, E. mitis, E. mivati, E. praecox and E. tenella. The particular content of the polyvalent vaccine, may be adjusted to contain different species depending on local conditions and the prevalence of particular species in certain areas. The most preferred vaccine contains all seven precocious species of Eimeria in the live attenuated form of sporulated oocysts.

The vaccine is administered to newly hatched chicks and preferably within one to two days of hatching. Following typical

- 3 -

husbandry techniques found in modern hatcheries, the vaccine is most preferably administered when the chicks are first processed in a process which administers other materials to the chick, such as other vaccines, for example Marek's disease, and perhaps vitamins and nutrients.

5 During the process the chicks are also usually debeaked. Thus, the instant vaccine finds its most preferred industrial utility when administered along with other administered materials such that the chicks are only handled once. However, on farms with a smaller scale of operations, it will not be uncommon to administer the instant vaccine
10 as a single administration without the concurrent administration of other materials and the simultaneous debeaking of the chicks.

The instant vaccine is generally administered as an aqueous suspension of the attenuated, precocious oocysts of Eimeria. To prevent the settling of the oocysts, the aqueous suspension can contain various
15 suspending agents, thickeners and preservatives to assure an even distribution of the oocysts throughout the suspension.

The dosages of the attenuated precocious oocysts ranges from about 5 to about 1000 oocysts per bird for each of the Eimeria species included in the vaccine since one species will not necessarily
20 confer immunity against a challenge from another species. A more preferred dosage will be from about 10 to 500 oocysts per species per bird, most preferably between 10 and 100 oocysts per species per bird. The precise number of each species present in the polyvalent vaccine will vary depending on the specific level of infectivity of each species
25 and it will be normal to have a vaccine with different numbers of each species to reflect such different characteristics of each species. Generally, the vaccine is administered in a total volume of from about 0.1 to 1.0 ml, preferably about 0.25 to 0.50 ml.

The techniques for the preparation of the attenuated
30 precocious strains of the different species of Eimeria are well known and utilize the procedure of serial passages of the coccidial strain through the host fowl or in embryonated eggs, or a combination of passages in fowl and eggs. Examples of such egg and fowl passages to

- 4 -

produce attenuated precocious strains is found in U.S. 4,438,097 to Shirley and U.S. 5,055,292 to McDonald, et al.

5 The vaccine discussed in McDonald, et al has been developed into a commercial product for chickens under the trademark of PARACOX. This is a combination of all seven principal species of Eimeria in an aqueous suspension. The product is labeled for administration to chickens from 5 to 9 days of age. Administration to chickens younger than 5 days is not recommended.

10 It will be appreciated by those skilled in the art that other strains of Eimeria exist which infect other animals than fowl, for example coccidiosis of cattle (E. zuernii, E. bovis) coccidiosis of swine (E. deblicieki, E. neodeblicieki, E. seabra and E. spinosa) as well as many other animals both avian and mammalian. Vaccines developed for such animals, based upon the use of attenuated precocious strains and
15 infective stages of the parasite, can be used to impart long-term immunity to the parasite.

The following example is provided in order to more fully demonstrate the operation of the instant invention. It should not be construed as limitative of the invention.

20

EXAMPLE 1

A trial was conducted to examine the immunizing ability of different levels of precocious, attenuated oocysts (PARACOX, Pitman-Moore) given to one-day-old birds. Two thousand one hundred seventy
25 six birds were assigned to 36 floor pens each containing 60 birds. In addition, 43 birds were maintained in batteries to be used as uninfected birds for challenge studies. The floor-pens were divided into four groups. Before placement in pens, birds of three groups were given
30 oocysts by gavage equivalent to 0.2, 1 or 5 times the recommended number of oocysts in the labeling for PARACOX. For the next 27 days the birds were reared in their pens as usual with feed and water provided without restriction. The fourth group of birds was fed a ration containing 60 ppm of salinomycin to suppress coccidial

- 5 -

development; from Day 27 on these birds received unmedicated feed. Temperature and humidity were maintained at comfortable levels appropriate for the age of the chickens. When the birds were 27 days old, birds were taken from each pen, weighed and placed in battery
5 cages. The birds remained on the same rations in cages as when on the floor. These birds were challenged with 10,000 sporulated oocysts of field strain of Eimeria tenella per bird after 24 hours in the cages. Feces were collected for Days 4 to 6 inclusive after challenge. The birds were weighed and necropsied at 8 days after challenge. The birds
10 remaining in the floor-pens were weighed and challenged with 30,000 sporulated oocysts of Eimeria maxima per bird on Day 28. Samples of feces were collected from each pen at intervals to monitor oocyst output from the challenged birds. A sample of birds were taken at 7 days after
15 challenge to determine lesion scores and all the birds were weighed.

The results of the study show that vaccinated birds were protected from the harmful effects of challenge (Tables 1 and 2) and that the birds showed significant weight gain and feed efficiency improvements over the unvaccinated birds. Table 3, 4 and 5 show the live weight of the birds, the average feed conversion per bird and the
20 average feed efficiency per bird.

25

30

- 6 -

TABLE 1

5 Summary of weight gains for 8 days after challenge, lesion scores and oocyst output from birds challenged with Eimeria tenella in cages after vaccination at one day of age with a mixture of oocysts (PARACOX).

Variable	Control	Salino ¹	Oocysts, (Paracox dose)		
			0.2X	1X	5X
Weight gain, g	539	549	558	609	550
Lesion score	2.6	1.9	0.2	0.4	0.2
Oocyst, 10 ⁶ /bird	75	29	<.1	.1	<.1

15 ¹Salinomycin included in ration at 60 ppm.

TABLE 2

20 Summary of weight gains for 7 days after challenge, lesion scores and oocyst output of birds challenged with Eimeria maxima after vaccination at one day of age with a mixture of oocysts (PARACOX).

Variable	Control ¹	Oocysts, (Paracox dose)		
		0.2X	1X	5X
Weight gain, g	279	435	465	463
Lesion score	2.1	1.0	0.9	1.1
Oocyst, 10 ⁶ /bird	16288	2440	3087	3775

30 ¹Birds fed ration with 60 ppm salinomycin until Day 27 and unmedicated ration thereafter.

- 7 -

TABLE 3

Treatments were replicated eight times. Average Live Weight, Feed Conversion and, Feed Efficiency of birds vaccinated with 0.20X, 1X or 5X level of PARACOX administered once by gavage on Day 0 and compared with unvaccinated birds fed 60 ppm Salinomycin up to Day 27 and unmedicated feed thereafter.

Day 28

10

VARIABLE	PARACOX			SALINOMYCIN
	0.2X	1X	5X	0-27 Days Unmedicated feed thereafter
Av. Live Weight (kg/bird)	1.085	1.085	1.080	1.077
Av. Feed Conversion g. of feed/ g. of weight gained	1.686	1.662	1.666	1.649
Av. of Feed Efficiency g. of weight gained /g. of feed	0.593	0.602	0.600	0.606

25

30

- 8 -

TABLE 4

Treatments were replicated eight times. Average Live Weight, Feed conversion and, Feed Efficiency of birds vaccinated with 0.20X, 1X or 5X level of PARACOX administered once by gavage on Day 0 and compared with unvaccinated birds fed 60 ppm Salinomycin up to Day 27 and unmedicated feed thereafter. Birds of all treatment groups were challenged on Day 28 with 30,000 sporulated oocysts of Eimeria maxima per bird via feed.

Day 35

VARIABLE	PARACOX			SALINOMYCIN
	0.2X	1X	5X	0-27 Days Unmedicated feed thereafter
Av. Live Weight (kg/bird)	1.520	1.550	1.543	1.356
Av. Feed Conversion g. of feed/ g. of weight gained	1.808	1.769	1.773	1.865
Av. of Feed Efficiency g. of weight gained /g. of feed	0.553	0.565	0.564	0.536

- 9 -

TABLE 5

Treatments were replicated eight times. Average Live Weight, Feed Conversion and, Feed Efficiency of birds vaccinated with 0.20X, 1X or 5X level of PARACOX administered once by gavage on Day 0 and compared with unvaccinated birds fed 60 ppm Salinomycin up to Day 27 and unmedicated feed thereafter. Birds of all treatment groups were challenged on Day 28 with 30,000 sporulated oocysts of Eimeria maxiuma per bird via feed.

<u>Day 42</u>				
VARIABLE	PARACOX			SALINOMYCIN
	0.2X	1X	5X	0-27 Days Unmedicated feed thereafter
Av. Live Weight (kg/bird)	2.080	2.093	2.096	1.924
Av. Feed Conversion g. of feed/ g. of weight gained	1.891	1.865	1.865	1.920
Av. of Feed Efficiency g. of weight gained /g. of feed	0.529	0.536	0.539	0.521

- 10 -

WHAT IS CLAIMED IS:

1. A method for inducing immunity to coccidiosis in animals which comprises administering to such animal a vaccine
5 composed of attenuated, precocious strains of sporulated oocysts of one or more species of Eimeria where the vaccine is administered orally upon birth or hatching of the fowl up to two days following the birth or hatching.
- 10 2. The method of Claim 1 where the vaccine is administered between birth or hatching of the animal and one day following the birth or hatching of the animal.
- 15 3. The method of Claim 1 where the animal being administered the vaccine is a fowl.
4. The method of Claim 2 where the fowl is chickens, turkeys, ducks or quail.
- 20 5. The method of Claim 3 where the fowl is chickens.
6. The method of Claim 1 where the vaccine is administered as an aqueous oral suspension of the sporulated oocysts.
- 25 7. The method of Claim 5 where the aqueous oral suspension also includes one or more suspending agents, thickeners or preservatives.
- 30 8. The method of Claim 1 where from 5 to 100 sporulated oocysts of each species of Eimeria are administered per animal.

- 11 -

9. The method of Claim 8 where from 10 to 500 sporulated oocysts of each species of Eimeria are administered per animal.

5 10. The method of Claim 9 where from 10 to 100 sporulated oocysts of each species of Eimeria are administered per animal.

10 11. The method of Claim 1 where the vaccine is composed of sporulated oocysts of one or more species of Eimeria selected from E. necatrix, E. acervulina, E. brunetti, E. maxima, E. mitis, E. praecox and E. tenella.

15 12. The method of Claim 11 where the vaccine is composed of sporulated oocysts of all of the species of Eimeria selected from E. necatrix, E. avervulina, E. brunetti, E. maxima, E. mitis, E. praecox and E. tenella.

20

25

30

INTERNATIONAL SEARCH REPORT

International application No.

PC 1/US94/00075

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) :A61K 39/00, 39/02; C12N 1/10, 1/36

US CL :424/88, 92; 435/245, 258

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/88, 92; 435/245, 258

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Abstract Avian Pathology, Volume 18, No. 2, issued 1989, Bedrnik et al, "Field Vaccination of Broilers Against Coccidiosis", Biodid znumber 88062625, pages 255-264, Abstract only.	1-12
X	Avian Pathology, Volume 15, issue 1986, Long et al, "Immunisation of Young Broiler Chickens with Low Level Infections of <u>Eimeria tenella</u> , <u>E. acervulina</u> or <u>E. maxima</u> ", pages 271-278, see entire document.	1-12
X	US, A, 4,438, 097 (Shirley) 20 March 1984, Abstract, col. 4, lines 30-53; col. 5, lines 30-35.	1-12

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be part of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

05 March 1994

Date of mailing of the international search report

MAR 28 1994

 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Facsimile No. NOT APPLICABLE

Authorized officer

H. F. SIDBERRY

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/00075

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Second Asian/Pacific Poultry Health Conference, Proceedings 112, issued 23-25 September 1988, Shirley, "Control of Coccidiosis with Vaccines", pages 129-157, see pages 131-133, 135-136.	1-12

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.